



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

seem so striking. While thus noticing a selected range in reference to dissemination, another peculiarity comes to mind, viz., that on these same hill sides, it seems to select a *zone* of growth. This zone commences about the middle of the hill and with this as a base extends about half way to the top; the plants being most thrifty and abundant on the southern exposures. There is certainly something peculiar in this limit of range, as the lower portions of the hills offer, to all appearances, as favorable conditions for growth as the upper. The same might also be remarked in reference to many localities back from the river.

However the special object of this note is to point out some peculiar effects of cultivation upon this plant. The specimens I have are from the garden of Mr. E. A. Rau, Bethlehem, Penn., and the peculiarities are that *the leaves are prominently 4 to 5 lobed—lobes acute and mucronate*. I have just examined some 40 specimens collected here with the following results: 80 per cent *entire to undulate*, and 20 per cent *distinctly toothed*; in no specimens that I have ever seen have the leaves appeared lobed. We further notice that the leaves of the cultivated *Jeffersonia* are 4 to 5 times as large as those found in the woods and the pod is also much larger and heavier. The flowers the writer has never seen, but would infer that they were also much larger. I take the liberty to append a note from Mr. Rau in reference to this plant. He says, "The *Jeffersonia* was taken from the garden of Mr. Jacob Wolle, of Bethlehem, Penn., and as the plant has been detected by Mr. Wolle near Narrowsville, Bucks county, along the Nockamixon rocks, this may be the source of the specimens of which we write." Mr. Rau further adds that however uncertain the length of time this has been cultivated by Mr. Wolle, he knows it to have been under cultivation more than 12 years. Doubtless many others have noticed variations as a result of cultivation and failed to give publicity to their observations. But it seems to me both interesting and profitable to notice such marked effects as the one just described.—A. H. YOUNG, Hanover, Ind.

NATURAL GRAFTING.—Some five miles north-west of Hanover I was surprised to find a good-sized tree of *Carpinus Americana*, or Water Beech, apparently growing on the trunk of *Fagus ferruginea*, or the common Beech. A closer examination showed that the trees had started close together, and instead of one crowding the other out, they had coalesced, their bark had become continuous, and below where they fork it is impossible to tell which is which. The fork looks like a regular branch, the Water Beech coming out at quite an angle, according to its usual habit.—J. M. C.

HABENARIA PERAMCENA, GRAY.—This showy Orchid has added much of beauty to our collections this summer. Orchids are exceedingly scarce in this range, and hence the advent of one in any quantity is most welcome to botanists. Now, the very interest and peculiar attractiveness of this plant this year lies in its abundance. It was first found here by the writer in July 1873, and has not been seen since by him, although I believe others have collected a very few specimens. Now this year it appears in considerable abundance both in the old locality and in many new ones, which in past years have been tramped over again and again without any finds. It is hardly reasonable to suppose seeds have been borne to six or seven localities, as many miles apart, during the past year, and, on the other hand, if these plants have lain dormant so long there is much interest in a rest of such long duration.—A. H. Y.

SPERMACOCE GLABRA, MICHX.—We are glad to be able to record the rediscovery of this desirable plant. In 1870 it was named in the list of Mr. A. H. Young, as growing in Jefferson county on the river bank. As no specimens were saved, the discovery was practically of no value, and after repeated searches along the river bank doubt began to arise in our minds whether we really were entitled to the plant or not. In 1874, in my incom-